

(19)  Canadian
Intellectual Property
Office

An Agency of
Industry Canada

Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

(11) **CA 2 013 188**

(13) **C**

(40) 14.03.2000

(43) 28.09.1990

(45) 14.03.2000

(12)

(21) 2 013 188

(51) Int. Cl.⁵:

**A61K 031/71, A61K 031/19,
A61K 031/405**

(22) 27.03.1990

(30) 07/329,451 US 28.03.1989

(72)

Fu, Cherng-Chyi Roger (US).
Lidgate, Deborah M. (US).

(73)

Syntex (U.S.A.) Inc.
3401 Hillview Avenue PALO ALTO XX (US).

(74)

DENNISON ASSOCIATES

(54) SYSTEME POUR CONSERVER LES PREPARATIONS OPHTALMIQUES

(54) PRESERVATIVE SYSTEM FOR OPHTHALMIC FORMULATIONS

(57)

Stable, clear, antimicrobially effective, ophthalmic formulations are disclosed which provide an antimicrobially effective preservative. The formulations include an ophthalmologically effective amount of a drug, which is a -COON group-containing non-steroidal anti-inflammatory drug (NSAID) in combination with an antibiotic drug, and a preservative system formed of a quaternary ammonium preservative and a nonionic polyoxyethylated octylphenol surfactant, all in an aqueous vehicle. The preservative system can be used with other formulations which require the preservative to be ophthalmologically acceptable and antimicrobially effective. These formulations are useful for treating diseases and/or conditions that are either caused by, associated with or accompanied by inflammatory processes, including, among others, glaucoma, cystoid macular edema, uveitis, diabetic retinopathy and conjunctivitis, or any trauma caused by eye surgery or eye injury. The ophthalmologically acceptable antibiotic is preferably tobramycin which has been found not to interfere with the rate of diffusion of the NSAID. The combination of the NSAID and antibiotic is particularly effective in simultaneously preventing and/or eliminating infection while preventing and/or eliminating inflammation.

OPIC
OFFICE DE LA PROPRIÉTÉ
INTELLECTUELLE DU CANADA



CIPO
CANADIAN INTELLECTUAL
PROPERTY OFFICE

(12)(19)(CA) **Brevet-Patent**

(11)(21)(C) **2,013,188**

(22) 1990/03/27

(43) 1990/09/28

(45) 2000/03/14

(72) Fu, Cherng-Chyi Roger, US

(72) Lidgate, Deborah M., US

(73) Syntex (U.S.A.) Inc., US

(51) Int.Cl.⁵ A61K 31/71, A61K 31/405, A61K 31/19

(30) 1989/03/28 (07/329,451) US

(54) **SYSTEME POUR CONSERVER LES PREPARATIONS
OPHTALMIQUES**

(54) **PRESERVATIVE SYSTEM FOR OPHTHALMIC
FORMULATIONS**

(57) Stable, clear, antimicrobially effective, ophthalmic formulations are disclosed which provide an antimicrobially effective preservative. The formulations include an ophthalmologically effective amount of a drug, which is a -COOH group-containing non-steroidal anti-inflammatory drug (NSAID) in combination with an antibiotic drug, and a preservative system formed of a quaternary ammonium preservative and a nonionic polyoxyethylated octylphenol surfactant, all in an aqueous vehicle. The preservative system can be used with other formulations which require the preservative to be ophthalmologically acceptable and antimicrobially effective. These formulations are useful for treating diseases and/or conditions that are either caused by, associated with or accompanied by inflammatory processes, including, among others, glaucoma, cystoid macular edema, uveitis, diabetic retinopathy and conjunctivitis, or any trauma caused by eye surgery or eye injury. The ophthalmologically acceptable antibiotic is preferably tobramycin which has been found not to interfere with the rate of diffusion of the NSAID. The combination of the NSAID and antibiotic is particularly effective in simultaneously preventing and/or eliminating infection while preventing and/or eliminating inflammation.



Industrie Canada Industry Canada

2013188

26280/2 FF

ABSTRACT OF THE DISCLOSURE

Stable, clear, antimicrobially effective, ophthalmic formulations are disclosed which provide an antimicrobially effective preservative. The formulations include an ophthalmologically effective amount of a drug, which is a -COOH group-containing non-steroidal anti-inflammatory drug (NSAID) in combination with an antibiotic drug, and a preservative system formed of a quaternary ammonium preservative and a nonionic polyoxyethylated octylphenol surfactant, all in an aqueous vehicle. The preservative system can be used with other formulations which require the preservative to be ophthalmologically acceptable and antimicrobially effective. These formulations are useful for treating diseases and/or conditions that are either caused by, associated with or accompanied by inflammatory processes, including, among others, glaucoma, cystoid macular edema, uveitis, diabetic retinopathy and conjunctivitis, or any trauma caused by eye surgery or eye injury. The ophthalmologically acceptable antibiotic is preferably tobramycin which has been found not to interfere with the rate of diffusion of the NSAID. The combination of the NSAID and antibiotic is particularly effective in simultaneously preventing and/or eliminating infection while preventing and/or eliminating inflammation.

30

35

3374M

26280/2 FF

2013188

5

-1-

PRESERVATIVE SYSTEM FOR OPHTHALMIC FORMULATIONSFIELD OF THE INVENTION

10 The present invention relates to improved
ophthalmic formulations which use an improved
preservative system comprising a quaternary ammonium
preservative and a stabilizing amount of a nonionic
polyoxyethylated octylphenol surfactant for ophthalmic
15 formulations of carboxyl ("COOH") group-containing
non-steroidal anti-inflammatory drugs ("NSAIDs") and
contain an ophthalmologically acceptable antibiotic,
preferably tobramycin. The invention also relates to
methods of using these formulations for treating
20 diseases and/or conditions that are either caused by,
associated with or accompanied by inflammatory
processes, including, among others, glaucoma, cystoid
macular edema, uveitis, diabetic retinopathy and
conjunctivitis, or any trauma caused by eye surgery or
25 eye injury. In addition, the formulation can be used to
treat bacterial infection.

BACKGROUND OF THE INVENTION

To be ophthalmologically acceptable, a formulation
30 must possess a number of characteristics to comply with

35

3374M

26280/2 FF

- 2 -

2013163

the general FDA requirements of being safe and effective. In that eyes are quite sensitive to pain, the formulation must be developed such that it causes little to no discomfort or stinging when administered.

5. This feature is particularly important to insure user compliance and important in that such formulations are often administered in order to relieve pain or inflammation. The ophthalmic use of NSAID compounds was disclosed in U.S. Patent No. 4,454,151, where NSAID compounds (such as those described in U.S. Patents 4,089,969; 4,232,038; 4,087,539 and 4,097,579) were exemplified in formulation with $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$, $\text{Na}_2\text{HPO}_4 \cdot \text{H}_2\text{O}$, NaCl , benzalkonium chloride ("BAC") and sterilized water. While the formulations described in the '151 patent were efficacious, a complex was found to form between the NSAID and the BAC.

Due to the formation of this complex, the formulations did not have the stability desired for shelf life in commercial applications. A reasonable minimum shelf life is at least about one year, representing sufficient time to package, ship, and store a formulation without having to replace expired stock too frequently.

An ophthalmic suspension containing a particular NSAID is disclosed in U.S. Patent No. 4,087,538 issued May 2, 1978. The suspension is aqueous based and can include benzalkonium chloride. Another ophthalmic formulation is disclosed in U.S. Patent No. 4,559,343 issued December 17, 1985. The formulation is aqueous based and includes an NSAID and a benzalkonium chloride preservative. A somewhat similar ophthalmic formulation is disclosed in U.S. Patent No. 4,607,038 issued August 19, 1986. This formulation includes a specific NSAID (pranoprofen) in an aqueous based formula with a known

3374M

26280/2 FF

2013188

- 3 -

preservative. U.S. Patent No. 4,474,751 issued October 2, 1984 discloses ophthalmic formulations which gel in the eye in order to increase the bioavailability of the drug. The '751 patent discloses a large number of different active ingredients and excipient material. When this disclosure is taken in view of the other patents discussed above and the publications cited in each of them, the vast number of different ways of creating an ophthalmic formulation becomes apparent. Although there may be a considerable number of possible formulations and variations thereof, only certain specific formulations will meet all the requirements for being ophthalmologically acceptable.

In general, an ophthalmic formulation contains an active compound and various ophthalmologically acceptable excipients, in the form of a solution, an ointment, a suspension, etc. In order for an excipient to be ophthalmologically acceptable, it must be non-irritating to the eye in combination with other excipients and an active ingredient. The excipients must not prevent the active ingredient from penetrating the blood-aqueous barrier and/or diffusing through the various ocular substructures to the site where it is pharmacologically active. The excipients can interact with each other or the active drug. Accordingly, care in formulating is required in that so many materials may be used. These materials generally include a tonicifier, a preservative, a surfactant, a buffering system, a chelating agent, a viscosity agent as well as other stabilizing agents. Ophthalmic formulations must be sterile and must be preserved with an effective anti-microbial agent.

Organo-mercurials (e.g., thimerosal, phenylmercuric acetate and phenylmercuric nitrate) have been used extensively as the preservative in ophthalmic

3374M

26280/2 FF

2013138

- 4 -

solutions. These compounds, however, pose difficulties due to potential mercury toxicity as well as poor chemical stability. Benzalkonium chloride, a quaternary ammonium compound, has been widely used in ophthalmic solutions, and is considered to be the preservative of choice. However, BAC has typically been considered to be incompatible with anionic drugs (e.g., salicylates or nitrates, etc.) and can be inactivated by surfactants.

Many NSAIDs (such as ketorolac, indomethacin, flurbiprofen, diclofenac, and suprofen) are being developed for ocular use because of their activity as anti-inflammatory agents as well as their ability to prevent cystoid macular edema.

These NSAIDs have proven to be incompatible with quaternary ammonium compounds such as BAC because they can form a complex with them, rendering the preservative less available to serve its function, as is the case with other ophthalmic drugs that contain a -COOH group. Thus, less preferred preservatives have been used in such ophthalmic formulations. For example, Ocufen Ophthalmic solution, the first NSAID (flurbiprofen) approved by the FDA for ophthalmic use, incorporates thimerosal (with EDTA) as its preservative system.

European published application 306,984 (published March 15, 1989) discloses a stable, clear, antimicrobially effective, ophthalmic formulation containing an NSAID and a preservative system formed of a quaternary ammonium preservative and a nonionic surfactant all in an aqueous vehicle. Although the formulations of this European laid-open application are useful in treating diseases that are either caused by, associated with, or accompanied by inflammatory processes, there is no indication that the formulations of the European laid-open application are effective in preventing or eliminating infection.

3374M

26280/2 FF

2013188

- 5 -

A need has continued to exist for a stable, clear, antimicrobial preservative effective ophthalmic formulation for NSAIDs with antibiotics using BAC as the preservative, and an improved preservative system for
5 -COOH group containing ophthalmic drugs to overcome both inflammation and infection.

SUMMARY OF THE INVENTION

A primary object of the invention is to describe and disclose a formulation containing an
10 ophthalmologically effective amount of an NSAID in combination with an antibiotic, a quaternary ammonium preservative and a stabilizing amount of a nonionic polyoxyethylated octylphenol surfactant, all in an aqueous vehicle.

15 A feature of the present invention is that it allows for the preparation of stable, i.e., clear and antimicrobially and antibiotically effective, NSAID-containing ophthalmic formulations without the need for an organo-mercurial preservative.

20 Another feature is that methods for treating ophthalmic diseases in mammals using the ophthalmic pharmaceutical formulations of the invention are provided.

An advantage of the present invention is that it is
25 useful in the treatment of diseases or conditions associated with or accompanied by inflammatory processes, including, among others, glaucoma, cystoid macular edema, uveitis, diabetic retinopathy and conjunctivitis, or any trauma caused by eye surgery or
30 eye injury and eliminating infection.

These and other objects, advantages and features of the present invention will become apparent to those persons skilled in the art upon reading the details of the composition, manufacture and usage as more fully set
35 forth below. Reference being made to the accompanying

3374M

26280/2 FF

- 6 -

2013132

general structural formulae forming a part hereof wherein like symbols refer to like molecular moieties throughout.

5 DETAILED DESCRIPTION AND PREFERRED EMBODIMENTS

Before the present compositions and processes for making and using such are disclosed and described, it is to be understood that this invention is not limited to the particular compositions, components or methods of use described as such compositions, components and methods may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting since the scope of the present invention will be limited only by the appended claims.

It must be noted that as used in this specification and the appended claims, the singular forms "a", "an" and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a pharmaceutically acceptable salt" includes mixtures of salts, references to "an NSAID" includes reference to mixtures of such NSAIDS, reference to "the method of administration" includes one or more different methods of administration known to those skilled in the art.

Definitions

As used herein, the term "NSAID" means an ophthalmologically acceptable carboxyl group containing non-steroidal anti-inflammatory drug. The NSAID's include, for example, flurbiprofen, ketorolac, diclofenac, indomethacin, suprofen, and the isomers, esters and pharmaceutically acceptable salts thereof.

As used herein, the term "q.s." means adding a

3374M

26280/2 FF

- 7 -

2013183

quantity sufficient to achieve a stated function, e.g., to bring a solution to the desired volume (i.e., 100%).

As used herein, the term "treatment" or "treating" means any treatment of a disease and/or condition in a mammal, including:

- (i) preventing the disease and/or condition, that is, causing the clinical symptoms of the disease not to develop;
- (ii) inhibiting the disease and/or condition, that is, arresting the development of clinical symptoms; and/or
- (iii) relieving the disease and/or condition, that is, causing the regression of clinical symptoms.

As used herein, the term "effective amount" means a dosage sufficient to provide treatment for the disease state being treated. This will vary depending on the patient, the disease and the treatment being effected.

As used herein, the term "antimicrobially effective" refers to the stability of the formulation prior to administration and means ability to withstand the U.S. Pharmacopia antimicrobial challenge put by a panel of microbes.

As used herein, the term "surfactant" means a nonionic surfactant, preferably ethoxylated octylphenol compounds as described below.

As used herein, the term "quarternary ammonium preservative" means a quarternary ammonium compound as described below.

As used herein, the term "stabilizing" means keeping a formulation clear and antimicrobially effective for its minimum reasonable shelf life, e.g., at least one year.

"Ketorolac tromethamine" shall mean the compound (\pm)-5-benzoyl-1,2-dihydro-3H-pyrrolo-[1,2-a]-pyrrole-1-carboxylic acid 2-amino-2-hydroxymethyl-1,3-propanediol

3374M


26280/2 FF

- 8 -

2013188

salt, also known as (\pm)-5-benzoyl-2,3-dihydro-1H-pyrrolizine-1-carboxylic acid with 2-amino-2-(hydroxymethyl)-1,3-propanediol (1:1) having the following structural formula (1)

5




10

"Tobramycin" shall mean the antibiotic produced by Streptomyces tenebrarius also known as O-3-amino-3-deoxy-a-D-glucopyranosyl-(1 β 6)-O-[2,6-diamino-2,3,6-trideoxy-a-D-ribo-hexopyranosyl-(1 β 4)]-2-deoxy-D-streptamine. Tobramycin is represented by the following structural formula II:

20

25

30



Tobramycin is a water soluble aminoglycosidic antibiotic having a broad spectrum of action against both gram negative and gram positive bacteria. Such

3374M

26280/2 FF

- 9 -

2013133

aminoglycosidic antibiotics are useful in treating ocular infections and are used prophylactically before and after ocular surgery.

Formulations

5 The formulations of the present invention include an NSAID active agent in an effective amount for ophthalmic treatment, an ophthalmologically acceptable antibiotic as a second active agent in an effective amount for ophthalmic treatment, a quaternary ammonium
10 preservative, a stabilizing amount of a nonionic polyoxyethylated octylphenol surfactant, optionally including other excipients such as a chelating agent, a tonicifier, a buffering system, a viscosity agent as well as other stabilizing agents.

15 The NSAID is preferably flurbiprofen, ketorolac, diclofenac, indomethacin, suprofen, and the isomers, esters, and pharmaceutically acceptable salts thereof. The antibiotic is preferably tobramycin.

 Ophthalmic solutions and suspensions typically
20 contain an aqueous vehicle rather than an oily vehicle. Ophthalmic formulations must be sterile, and if intended for multiple dosing regimens, must be antimicrobially effective for their minimum reasonable shelf life, e.g.,
25 at least one year, and preferably two to three years or more. The ingredients used in the formulations of the present invention are typically commercially available or can be made by methods readily known to those skilled in the art.

 Pharmaceutical ophthalmic formulations typically
30 contain an effective amount, e.g., 0.001% to 10% wt/vol., preferably 0.002% to 5% wt/vol, most preferably 0.005% to 1% of an active ingredient (e.g., the NSAID of the present invention). The amount of active ingredient will vary with the particular formulation and the
35 disease state for which it is intended. The total

3374M

26280/2 FF

- 10 -

2013188

concentration of solutes should be such that, if possible, the resulting solution is isotonic with the lacrimal fluid (though this is not absolutely necessary) and has a pH in the range of 6 to 8.

- 5 The formulations of the present invention are prepared as solutions incorporating the above-described ingredients within the following approximate ranges:

	<u>Ingredient</u>	<u>Amount</u>
	Active Agent*	0.001% to 10.0% wt/vol.;
10	Preservative	0.001% to 1.0% wt/vol.;
	Surfactant	0.001% to 1.0% wt/vol.;
	Other Excipients	0% to 10.0% wt/vol.; and
	Purified Water	q.s. to 100%.

- 15 *The active agent is the NSAID in combination with the antiobiotic.

Optional other excipients, such as a chelating agent and a tonicifier, are used in the following approximate proportions:

	<u>Ingredient</u>	<u>Amount</u>
20	Chelating agent	0.01% to 1.0%wt/vol.;
	Tonicifier	q.s. to achieve isotonicity with lacrimal fluid; and
25	1N NaOH or 1N HCl	q.s. to adjust pH to 6.0 to 8.0.

30

35

3374M

26280/2 FF

- 11 -

2013189

In a preferred ophthalmic NSAID solution, the ingredients are combined in the following proportions:

	<u>Ingredient</u>	<u>Amount</u>
	NSAID	0.50% wt/vol.;
5	Antibiotic	0.30% wt/vol.;
	BAC	0.02% wt/vol.;
	(50% aq. soln.)	
	Octoxynol 40	0.01% wt/vol.;
	(70% aq. soln.)	
10	EDTA Na ₂	0.10% wt/vol.;
	NaCl/ boric acid/	q.s. for isotonicity with
	Na borate	lacrimal fluid;
	1N NaOH or 1N HCl	q.s. to adjust pH to
		7.4±0.4; and
15	Purified Water	q.s. to 100%.

The invention relates primarily to formulations having as the active agent ophthalmologically acceptable drugs (including the esters and pharmaceutically acceptable salts thereof) that can form a complex with a quaternary ammonium compound, particularly carboxyl group-containing NSAIDs.

NSAIDs useful in the practice of this invention include, for example, ketorolac (and the other compounds described as being ophthalmologically effective in U.S. Patent No. 4,454,151 to Waterbury, issued June 12, 1984, the pertinent portions of which are incorporated herein by reference), indomethacin, flurbiprofen sodium, diclofenac, and suprofen, including the esters and pharmaceutically acceptable salts thereof.

In addition to the NSAID there is another active ingredient in the form of an ophthalmologically acceptable antibiotic, preferably tobramycin. The antibiotic is present in an effective amount for ophthalmic treatment. The antibiotic tobramycin does not interfere with the corneal permeability of the NSAID.

3374M

26280/2.FF

- 12 -

2013138

Preservatives useful in the formulations of the present invention include quaternary ammonium compounds, such as cetyltrimethylammonium bromide, cetylpyridinium chloride and preferably, benzalkonium chloride.

- 5 The nonionic surfactants useful in the formulations of the present invention are preferably polyoxyethylated octylphenol surfactants including polyoxyethylene hydrogenated vegetable oils, such as polyethylene 60 hydrogenated castor oil, manufactured and sold by Kao Corp. of Japan under the trade name Emanon CH-60, and
10 preferably ethoxylated octylphenol compounds, such as Octoxynol 10 and most preferably Octoxynol 40, manufactured and sold by GAF under the trade name Igepal CA897 (a 70% aqueous solution of Octoxynol 40).
15 Octoxynol 40 is a nonionic polymeric surfactant material. More specifically, it is a nonionic polyoxyethylated octylphenol surfactant material sold commercially by GAF.

- Among the optional excipients, the chelating agents
20 useful in the formulations of the present invention include 8-hydroxyquinoline sulfate, citric acid, and preferably disodium edetate. Under certain conditions, the chelating agent may also enhance the anti-microbial effect due to its ability to render essential metal ions
25 unavailable to the microbes.

 Buffering systems optionally useful in the formulations of the present invention are based on, for example, citrate, borate, or phosphate.

- Tonicifiers optionally useful in the formulations
30 of the present invention include dextrose, potassium chloride and/or sodium chloride, preferably sodium chloride.

- Viscosity agents optionally useful in the formulations of the present invention include the
35 cellulose derivatives such as hydroxypropylmethyl

3374M

26280/2 FF

- 13 -

2013188

cellulose, sodium carboxymethylcellulose, and hydroxyethylcellulose.

Other optional excipients useful in the formulations of the present invention include stabilizing agents such as antioxidants, e.g., sodium metabisulfate and ascorbic acid, depending on the NSAID used.

These formulations are prepared by dissolving the solutes (e.g., the NSAID, the preservative, the surfactant, the chelating agent, and the buffering agent) in a suitable quantity of water, adjusting the pH to about 6 to 8, preferably 6.8 to 8.0 and most preferably 7.4, making a final volume adjustment to 100% with additional water, and sterilizing the preparation using any suitable method known to those in the art.

Ophthalmic formulations incorporating the preservative system of the invention are physically stable (i.e., remain clear) and functionally stable (i.e., remain antimicrobially effective) for at least the minimum reasonable shelf life of such products. The inclusion of an antibiotic in the formulation does not effect the rate of diffusion of the NSAID.

Preferred Formulations

The preferred ophthalmic formulation of the invention includes a NSAID active agent in an effective amount for ophthalmic treatment and an antimicrobially effective amount of the above-described preferred preservative system.

The preferred preservative of the invention is benzalkonium chloride.

The preferred surfactant of the invention is Octoxynol 40, especially when combined with benzalkonium chloride.

3374M

26280/2 FF

2013183

- 14 -

The preferred chelating agent of the invention is disodium edetate, especially when combined with benzalkonium chloride and Octoxynol 40.

The preferred antibiotic is one which does not interfere with the corneal permeability of the NSAID. Tobramycin is a preferred antibiotic.

The preferred ophthalmic solutions of the invention include a NSAID, benzalkonium chloride, Octoxynol 40 and disodium edetate and, as a second active agent, tobramycin.

A preferred ophthalmic NSAID/antibiotic solution has the following formulation:

	<u>Ingredient</u>	<u>Amount</u>
	NSAID	0.50% wt/vol.
15	antibiotic	0.30% wt/vol.
	BAC	0.02% wt/vol.
	(50% aq. soln.)	
	Octoxynol 40	0.01% wt/vol.
	(70% aq. soln.)	
20	EDTA Na ₂	0.10% wt/vol.
	(NaCl/boric acid/ Na borate)	q.s. for isotonicity with lacrimal fluid
	1N NaOH or 1N HCl	q.s. to adjust pH to 7.4±0.4
25	Purified Water	q.s. to 100%

Most preferred is the ophthalmic solution according to the above formulations is wherein the NSAID is Ketorolac Tromethamine and when the antibiotic is present it is tobramycin.

Utility and Administration

This invention is directed to NSAID ophthalmic formulations and a method useful for treating ophthalmic diseases in mammals. These diseases are either caused by, associated with or accompanied by inflammatory

3374M

26280/2 FF

2010168

- 15 -

processes, including, among others, glaucoma, cystoid macular edema, uveitis, diabetic retinopathy and conjunctivitis, or any trauma caused by eye surgery or eye injury.

- 5 The method of this invention is both curative and preventative. Where applied, for example, pre-surgically or immediately post-traumatically, i.e. before inflammation develops, it prevents development of inflammation. When applied directly to the eye
10 suffering from any of the named ophthalmic diseases, it suppresses already developed inflammatory processes.

- The formulation of the invention includes an antibiotic such as tobramycin, providing antibacterial properties useful in eliminating and/or preventing a
15 bacterial infection.

- Ophthalmic formulations are typically administered by topical application to the eyelids or for instillation into the space (cul-de-sac) between the eyeball and the eyelids, by topically applied ophthalmic
20 solutions, suspensions or ointments, or by subconjunctival injection.

- The dosage level will, of course, depend on the concentration of the drops, the condition of the subject and the individual magnitude of responses to treatment.
25 However, typical dosage ranges might be about 2 to 10 drops of solution of active ingredient per day wherein the solution includes 0.5 wt/vol.% of Ketorolac trimethamine and 0.3 wt/vol.% of tobramycin.

- For a more detailed discussion of ophthalmic
30 formulations, their preparation and administration, see Remington's Pharmaceutical Sciences, 15th Ed., pages 1489-1504, (1975).

Testing

- 35 Ophthalmic formulations such as the solutions of

3374M

26280/2 FF

2613183

- 16 -

the present invention are typically tested for physical stability, chemical stability, and preservative efficacy, both when they are first manufactured and after a fixed period of time (e.g., after two years).

- 5 They are generally considered to be safe and clinically acceptable if proven to be well tolerated in the eye.

- Physical stability is determined by observation of a solution after expiration of a fixed period of time. A solution is considered to be physically stable if its
10 appearance (e.g., color and clarity) does not change and if the pH remains constant, within acceptable limits. Chemical stability involves a routine chemical analysis of the solution, to be sure that its active
ingredient(s), preservatives and the excipients have not
15 changed after a fixed period of time.

- Preservative efficacy of the formulation prior to administration is tested by the procedure described in the U.S. Pharmacopia Compendiary, whereby a solution is challenged with a panel of microbes and a determination
20 is made as to whether a given microbe survives in it.

EXAMPLES

- The following examples are given to enable those
25 skilled in the art to more clearly understand and to practice the present invention. They should not be considered as a limitation on the scope of the invention, but merely as being illustrative and representative thereof.

30

EXAMPLE 1

- This example illustrates the preparation of a representative pharmaceutical formulation for ophthalmic administration containing the NSAID Ketorolac
35

3374M

26280/2 FF

2013133

- 17 -

Tromethamine and the antibiotic tobramycin.

	<u>Ingredient</u>	<u>Amount</u>
	ketorolac tromethamine	0.50% wt/vol.
5	tobramycin	0.30% wt/vol.
	BAC	0.02% wt/vol.
	(50% aq. soln.)	
	Octoxynol 40	0.01% wt/vol.
	(70% aq. soln.)	
10	EDTA Na ₂	0.10% wt/vol.
	NaCl	0.18% wt/vol.
	Boric Acid	0.9% wt/vol.
	Na Borate	0.45% wt/vol.

15 The above ingredients are mixed, adding purified water until they are dissolved, the pH is adjusted to 7.4±0.4 and the balance of the formulation is made up with purified water, adding a quantity sufficient to make 100% volume. The solution is then sterilized.

20 Other NSAIDs, such as those described above, can be used as the active compound in the preparation of the formulation of this example.

EXAMPLE 2

25 This example illustrates the preparation of a general pharmaceutical formulation for ophthalmic

30

35

3374M

26280/2 FF

2013183

- 18 -

administration containing an NSAID and an antibiotic.

	<u>Ingredient</u>	<u>Amount</u>
5	NSAID	0.50% wt/vol.
	antibiotic	0.3% wt/vol.
	BAC	0.01% wt/vol.
	(50% aq. soln.)	
	Octoxynol 40	0.02% wt/vol.
	(70% aq. soln.)	
10	EDTA Na ₂	0.20% wt/vol.
	NaCl	0.18% wt/vol.
	Boric Acid	0.9% wt/vol.
	Na Borate	0.45% wt/vol.

15

EXAMPLE 3

This example illustrates the preparation of a representative pharmaceutical formulation for ophthalmic administration containing the NSAID ketorolac tromethamine and tobramycin.

	<u>Ingredient</u>	<u>Amount</u>
25	ketorolac tromethamine	0.50% wt/vol.
	tobramycin	0.30% wt/vol.
	BAC	0.01% wt/vol.
	(50% aq. soln.)	
	Octoxynol 40	0.01% wt/vol.
	(70% aq. soln.)	
30	EDTA Na ₂	0.20% wt/vol.
	NaCl	0.18% wt/vol.
	Boric Acid	0.9% wt/vol.
	Na Borate	0.45% wt/vol.

Other NSAIDs, such as those described above, can be used as the active compound in the preparation of the

3374M

26280/2 FF

2013130

- 19 -

formulation of any of these examples.

EXAMPLE 4

This example illustrates the preparation of a
5 representative pharmaceutical formulation for ophthalmic
administration containing the NSAID flurbiprofen sodium.

	<u>Ingredient</u>	<u>Amount</u>
	Flurbiprofen Sodium	0.03% wt/vol.
10	BAC	0.02% wt/vol.
	(50% aq. soln.)	
	Octoxynol 40	0.01% wt/vol.
	(70% aq. soln.)	
	EDTA Na ₂	0.10% wt/vol.
15	NaCl	0.18% wt/vol.
	Boric Acid	0.9% wt/vol.
	Na Borate	0.45% wt/vol.

The above ingredients are mixed, adding purified
20 water until they are dissolved, the pH is adjusted to
7.4±0.4 and the balance of the formulation is made up
with purified water, adding a quantity sufficient to make
100% volume. The solution is then sterilized.

Other ophthalmic drugs and NSAIDs, such as those
25 described above, can be used as the active compound in
the preparation of the formulation of this example.

EXAMPLE 5

Physical stability of the formulations of the
30 present invention is measured by preparing clear
formulations, e.g., according to the foregoing Examples,
sealing them in sterilized containers, and observing the
clarity of the solution after a period of one month and
again after five months. Solutions that remain clear are
35 considered stable in this procedure.

3374M

26280/2 FF

2013188

- 20 -

The formulations of the present invention have proven to be stable when tested in accordance with the above procedure. Formulations using surfactants other than the nonionic surfactants of the invention did not remain clear and were not stable.

Preservative efficacy of the formulations of the present invention is measured by preparing formulations, e.g., according to the foregoing Examples, and subjecting them to the U.S. Pharmacopia antimicrobial challenge.

The formulations of the present invention demonstrate preservative efficacy when tested in accordance with the above procedure.

Formulations of the present invention are freely flowable liquids which can be administered directly to the eye using a conventional means such as eyedroppers. The amount of active ingredient administered will vary with the individual and/or the type of disease or condition being treated. The NSAID's such as ketorolac and antibiotics such as tobramycin are generally administered in an amount of about 1 to 2 drops per eye with drops containing about 25 microliters of formulation. The drops are generally administered 3 to 4 times per day.

25

EXAMPLE 6

In vitro rabbit corneal penetration of ketorolac was evaluated in the presence of tobramycin to determine if tobramycin alters penetration of ketorolac through rabbit corneas. Two sets of studies were performed to evaluate tobramycin's effect on ketorolac penetration.

Apparatus - A modified Franz diffusion cell consisting of an 8.0 ml glass receptor cell along with a teflon donor cell were used for the penetration experiments. A side arm allowed sampling of the receptor

3374M

26280/2 FF

- 21 -

2013120

- phase. The donor cell was recessed to accommodate corneal curvature. A 0.3 ml volume of donor solution was placed on the epithelial side of the cornea, and evaporation of this donor solution was diminished by
- 5 sealing a glass coverslip over the opening of the donor cell with silicon grease. To ensure corneal curvature throughout the course of the experiment, a 1.0 ml latex bulb was placed over the sampling port of the glass diffusion cell. By so doing, enough pressure was exerted
- 10 under the cornea to maintain a curved, wrinkle-free membrane. Water at 37° C was circulated through the water jacket surrounding the receptor cell. A magnetic stir bar placed in the bottom of the receptor cell maintained homogeneity within the receptor solution.
- 15 Cornea Preparation - New Zealand white rabbits weighing 3.5 to 4 kg were used for the studies. Rabbits were sacrificed by rapid injection of 1.25 ml/kg of T-61 Euthanasia Solution (American Hoechst Corp. Animal Health Division, Somerville, NJ) into a marginal ear vein. The
- 20 cornea were carefully removed along with 2-4 mm of surrounding scleral tissue then placed in a buffer containing: 0.57% sodium chloride, 0.361% sodium bicarbonate, 0.04% potassium chloride, 0.023% potassium phosphate dibasic, 0.007% magnesium sulfate, 0.08%
- 25 calcium chloride, and 0.133% adenosine in water, adjusted to pH 7.4. This buffer was used as receptor solution for all studies; its selection was based on the ability to maintain corneal integrity throughout the diffusion studies.
- 30 Experimental Procedure - A fresh cornea was placed between the top and bottom of the teflon donor cell; this unit was then clamped onto the glass receptor cell. The receptor cell was filled with sterile, degassed buffer solution; all air bubbles were expelled from beneath the
- 35 cornea by inverting the entire diffusion cell and

3374M

26280/2 FF

2018138

- 22 -

allowing bubbles to travel out the sampling port. After donor solution was placed on the cornea, a 0.3 ml sample of receptor solution was collected at the following time points: 15, 30, 45, 60 and 120 minutes. The 0.3 ml aliquot was replaced at each time point with fresh buffer solution.

Preparation of Test Solutions - 1. To determine ketorolac corneal diffusion in the presence of tobramycin, and to determine a dose effect, a saline vehicle was utilized to avoid potential complications by excipients. The following solutions were isotonic and prepared at pH 7.4: (a) 0.5% ketorolac tromethamine, 0.79% sodium chloride, purified water; (b) solution (a) with 0.15% tobramycin; (c) solution (a) with 0.30% tobramycin; and (d) solution (a) with 0.60% tobramycin.

2. To evaluate whether 0.30% tobramycin (a clinically acceptable and efficacious concentration) has an effect on ketorolac corneal diffusion when administered in a more complex vehicle, an isotonic solution at pH 7.4 was made which contained the following: (a) 0.5% ketorolac tromethamine, 0.79% sodium chloride, edetate disodium, benzalkonium chloride, purified water; (b) solution (a) with 0.30% tobramycin.

¹⁴C-glycerol Penetration - To monitor corneal integrity throughout the course of the permeability studies, ¹⁴C-glycerol penetration was evaluated (¹⁴C-glycerol 15.76 mCi/mmol was obtained from NEN with a radiochemical purity of 98%). Nonionized ¹⁴C-glycerol was incorporated into selected test solutions (1a and d, above). For controls, two additional isotonic test solutions were made at pH 7.4: (1) phosphate buffered saline; (2) 0.6% tobramycin in phosphate buffered saline. To a 2.0 ml aliquot of each test solution, 10 µl of ¹⁴C-glycerol was added. At designated time intervals, 0.3 ml of receptor solution

3374M

26280/2 FF

2013188

- 23 -

was sampled for scintillation counting (Beckman model LS 8100).

Analytical Methods - 1. Quantitation of ketorolac was performed by HPLC. The mobile phase was composed of
5 methanol, water and glacial acetic acid (65:34:1). The equipment included: a Spectra-Physics 8440 UV/Vis detector; a Spectra-Physics 4270 integrator; a Spectra-Physics 8700 solvent delivery system; a Dynatech autosampler; and a Whatman Partisil ODS 3, 10 micron
10 column. The mobile phase flow rate was 1.0 ml/min; the sample injection volume was 50 µl; and the absorbance wavelength was 254 nm. A 100 µl aliquot of each sample was diluted with 150 µl of mobile phase.

2. Quantitation of tobramycin was performed using
15 the Syva EMIT tobramycin assay kit. The assay is an enzyme immunoassay intended to quantitatively analyze tobramycin in human serum or plasma; the limit of detection is 1.0 µg/ml. The assay is based on competition for antibody sites between free drug in
20 sample and drug labeled with glucose-6-phosphate dehydrogenase (G-6-P-DH). Since G-6-P-DH activity decreases upon binding with antibody, tobramycin concentration can be measured in terms of enzyme activity. Active enzyme converts oxidized nicotinamide
25 adenine dinucleotide (NAD) to NADH. This conversion results in an absorbance change that is measured spectrophotometrically.

Each experiment was performed with matched controls; that is, from a single rabbit, one cornea was treated
30 with a ketorolac (control) solution, and the other cornea was treated with the ketorolac and tobramycin solution. Each test solution containing tobramycin was evaluated in triplicate. For the study using the simple isotonic vehicle, data for nine control corneas were generated.
35 Since these were control cornea, each is from a different

3374M

26280/2 FF

2013188

- 24 -

rabbit; hence, the deviation shown at each time point gives an indication of both the biological as well as experimental deviation inherent to this type of study.

An indication of corneal integrity throughout the course of these studies was determined by penetration of ^{14}C -glycerol. Changes in the permeability profile of ^{14}C -glycerol can be attributed to corneal alteration or damage. Select vehicles were chosen to evaluate whether corneal damage could be attributed to a particular compound or combination. With phosphate buffered saline serving as control, a two or three-fold increase in ^{14}C -glycerol penetration would indicate substantial corneal alteration. Table I shows that ^{14}C -glycerol penetration in a solution containing ketorolac tromethamine, or 0.6% tobramycin, or their combination, does not differ from its penetration in buffer alone. These results suggest that corneal integrity is not altered by ketorolac tromethamine or tobramycin.

20

TABLE I

	Preparation	Percent of Initial Counts per Minute	
		at 60 min	at 120 min
25	Phosphate Buffered Saline	2.10	7.36
	Ketorolac tromethamine in Saline	2.47	8.60
	Tobramycin (0.6%) in Phosphate buffered saline	1.83	7.08
30	Ketorolac tromethamine and Tobramycin (0.6%) in Saline	2.01	6.03

The average total milligrams of ketorolac penetrating the cornea at each time point for the simple solutions containing ketorolac alone and solutions

35 3374M

26280/2 FF

2013188

- 25 -

containing either 0.15%, 0.30% or 0.60% tobramycin, respectively, were compared. In all cases, the solutions containing tobramycin were equivalent to the control solution.

- 5 A comparison of the average total milligrams of ketorolac penetrating the cornea at each time point for the ophthalmic formulation with and without 0.30% tobramycin was made. Again, the test solution and the control solution were equivalent. Studies with the
- 10 formulation demonstrated that after 60 minutes, there occurs a two to three fold increase in ketorolac diffusion, that is, enhanced penetration.

- While the present invention has been described with
- 15 reference to the specific embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. In addition, many modifications may be made
- 20 to adapt a particular situation, material, composition of matter, process, process step or steps, to the objective, spirit and scope of the present invention. All such modifications are intended to be within the scope of the claims appended hereto.

25

30

35

3374M

26280/2 FF

2013188

- 26 -

WHAT IS CLAIMED IS:

1. An ophthalmologically acceptable non-steroidal anti-inflammatory drug formulation, comprising:
 - 5 an ophthalmologically acceptable non-steroidal anti-inflammatory carboxyl group-containing drug in an effective amount for ophthalmic treatment;
 - an ophthalmologically acceptable antibiotic in an effective amount for ophthalmic treatment;
 - 10 a quaternary ammonium preservative;
 - a stabilizing amount of a nonionic polyoxyethylated octylphenol surfactant; and
 - an aqueous vehicle.
- 15 2. The ophthalmologically acceptable non-steroidal anti-inflammatory drug formulation of Claim 1 wherein said quaternary ammonium preservative is benzalkonium chloride.
- 20 3. The ophthalmologically acceptable non-steroidal anti-inflammatory drug formulation of Claim 1 wherein said nonionic polyoxyethylated octylphenol surfactant is Octoxynol 40 and the antibiotic is tobramycin.
- 25 4. The ophthalmologically acceptable non-steroidal anti-inflammatory drug formulation of Claim 1 including disodium edetate.
- 30 5. The ophthalmologically acceptable non-steroidal anti-inflammatory drug formulation of Claim 1 wherein said ophthalmologically acceptable non-steroidal anti-inflammatory carboxyl group-containing drug is selected from the group: ketorolac, indomethacin, flurbiprofen, diclofenac, and suprofen.

35

3374M

26280/2 FF

- 27 -

2013188

6. The ophthalmologically acceptable non-steroidal anti-inflammatory drug formulation of Claim 5 wherein said ophthalmologically acceptable non-steroidal anti-inflammatory carboxyl group-containing drug is ketorolac tromethamine.

7. The ophthalmologically acceptable non-steroidal anti-inflammatory drug formulation of Claim 1 comprising:

10	NSAID	0.001% to 10.0% wt/vol.;
	Antibiotic	0.001% to 10.0% wt/vol.;
	Preservative	0.001% to 1.0% wt/vol.;
	Surfactant	0.001% to 1.0% wt/vol.;
		and
15	Purified Water	q.s. to 100%.

8. The ophthalmologically acceptable non-steroidal anti-inflammatory drug formulation of Claim 7 comprising:

20	ketorolac tromethamine	0.001% to 10.0% wt/vol.;
	tobramycin	0.001% to 10.0% wt/vol.;
	Preservative	0.001% to 1.0% wt/vol.;
	Surfactant	0.001% to 1.0% wt/vol.;
	Purified Water	q.s. to 100%.

25

9. The ophthalmologically acceptable non-steroidal anti-inflammatory drug formulation of Claim 7 wherein said preservative is benzalkonium chloride, and the surfactant is Octoxynol 40.

30

10. The ophthalmologically acceptable non-steroidal anti-inflammatory drug formulation of

35

3374M

26280/2 FF

2013188

- 28 -

Claim 8, further comprising:

Chelating agent 0.01% to 1.0% wt/vol.;
Tonicifier q.s. to achieve isotonicity
5 with lacrimal fluid; and
1N NaOH or 1N HCl q.s. to adjust pH to
7.4%0.4.

11. The opthalmologically acceptable
10 non-steroidal anti-inflammatory drug formulation of
Claim 9 comprising:

ketorolac tromethamine 0.50% wt/vol.;
Tobramycin 0.30% wt/vol.;
BAC 0.02% wt/vol.;
15 (50% aq. soln.)
Octoxynol 40 0.01% wt/vol.;
(70% aq. soln.)
EDTA Na₂ 0.10% wt/vol.;
NaCl 0.18% wt/vol.;
20 Boric Acid 0.9% wt/vol.
Na Borate 0.45% wt/vol.
1N NaOH or 1N HCl q.s. to adjust pH to
7.4%0.4; and
Purified Water q.s. to 100%.
25

12. The use of a formulation comprising: an
ophthalmologically acceptable non-steroidal anti-
inflammatory carboxyl group-containing drug in an effective
amount for ophthalmic treatment, an antibiotic in an
effective amount for ophthalmic treatment, a quaternary
ammonium preservative, a stabilizing amount of a nonionic
polyoxyethylated octylphenol surfactant, and an aqueous

3374M

26280/2 FF

B

2013188

- 29 -

vehicle for treating ophthalmic disease in a mammal suffering therewith.

13. The use of Claim 12 wherein said preservative is benzalkonium chloride and said surfactant is Octoxynol 40.

14. The use of Claim 12 wherein said ophthalmologically acceptable non-steroidal anti-inflammatory carboxyl group-containing drug is selected from the group: ketorolac, indomethacin, flurbiprofen, diclofenac, and suprofen.

15. The use of Claim 12 wherein said ophthalmologically acceptable non-steroidal anti-inflammatory carboxyl group-containing drug is Ketorolac Tromethamine and the antibiotic is tobramycin.

16. The use of Claim 15 wherein said ophthalmologically acceptable non-steroidal anti-inflammatory drug formulation comprises:

25	ketorolac tromethamine	0.50% wt/vol.;
	Tobramycin	0.30% wt/vol.;
	BAC	0.01% wt/vol.;
	(50% aq. soln.)	
	Octoxynol 40	0.01% wt/vol.;
	(70% aq. soln.)	
30	EDTA Na ₂	0.10% wt/vol.;
	NaCl	0.18% wt/vol.;
	Boric Acid	0.9% wt/vol.
	Na Borate	0.45% wt/vol.
	1N NaOH or 1N HCl	to adjust pH to
35		7.4±0.4; and
	Purified Water	q.s. to 100%.

3374M

26280/2 FF

B